



MARKSCHEME

May 2010

MATHEMATICS

Higher Level

Paper 1

Samples to team leaders	June 7 2010
Everything (marks, scripts etc) to IB Cardiff	June 14 2010

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of IB Cardiff.*

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Write the marks in red on candidates' scripts, in the right hand margin.

- Show the **breakdown** of individual marks awarded using the abbreviations **MI**, **AI**, etc.
- Write down the total for each **question** (at the end of the question) and **circle** it.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **AI**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, e.g. **MIAI**, this usually means **MI** for an **attempt** to use an appropriate method (e.g. substitution into a formula) and **AI** for using the **correct** values.
- Where the markscheme specifies (**M2**), **N3**, etc., do **not** split the marks.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 N marks

*Award N marks for **correct** answers where there is **no** working.*

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

4 Implied marks

*Implied marks appear in **brackets e.g. (M1)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.*

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 Follow through marks

*Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.*

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (e.g. $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 Mis-read

*If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). Apply a **MR** penalty of 1 mark to that question. Award the marks as usual and then write $-1(\mathbf{MR})$ next to the total. Subtract 1 mark from the total for the question. A candidate should be penalized only once for a particular mis-read.*

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).

7 Discretionary marks (*d*)

*An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. The mark should be labelled (**d**) and a brief **note** written next to the mark explaining this decision.*

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2\sin(5x - 3)$, the markscheme gives:

$$f'(x) = 2\cos(5x - 3) \cdot 5 = 10\cos(5x - 3) \quad \mathbf{AI}$$

Award **AI** for $2\cos(5x - 3) \cdot 5$, even if $10\cos(5x - 3)$ is not seen.

10 Accuracy of Answers

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy.

- **Rounding errors:** only applies to final answers not to intermediate steps.
- **Level of accuracy:** when this is not specified in the question the general rule applies: *unless otherwise stated in the question all numerical answers must be given exactly or correct to three significant figures.*

Candidates should be penalized **once only IN THE PAPER** for an accuracy error (**AP**). Award the marks as usual then write (**AP**) against the answer. On the **front** cover write $-1(\mathbf{AP})$. Deduct 1 mark from the total for the paper, not the question.

- If a final correct answer is incorrectly rounded, apply the **AP**.
- If the level of accuracy is not specified in the question, apply the **AP** for correct answers not given to three significant figures.

If there is no working shown, and answers are given to the correct two significant figures, apply the **AP**. However, do **not** accept answers to one significant figure without working.

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

SECTION A

1. using the factor theorem or long division *MI*
 $-A + B - 1 + 6 = 0 \Rightarrow A - B = 5$ *AI*
 $8A + 4B + 2 + 6 = 0 \Rightarrow 2A + B = -2$ *AI*
 $3A = 3 \Rightarrow A = 1$ *AI*
 $B = -4$ *AI* N3

Note: Award *MIA0A0AIAI* for using $(x - 3)$ as the third factor, without justification that the leading coefficient is 1.

[5 marks]

2. $g(x) = 0$ or 3 *(MI)(AI)*
 $x = -1$ or 4 or 1 or 2 *AIAI*

Notes: Award *AIAI* for all four correct values,
AIA0 for two or three correct values,
A0A0 for less than two correct values.

Award *MI* and corresponding *A* marks for correct attempt to find expressions for f and g .

[4 marks]

3. (a) for using normal vectors *(MI)*
 $\begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} = 1 - 1 = 0$ *MIAI*
hence the two planes are perpendicular *AG*

(b) **METHOD 1**

EITHER

$$\begin{vmatrix} i & j & k \\ 1 & 2 & -1 \\ 1 & 0 & 1 \end{vmatrix} = 2i - 2j - 2k$$
MIAI

OR

if $\begin{pmatrix} a \\ b \\ c \end{pmatrix}$ is normal to π_3 , then

$$a + 2b - c = 0 \text{ and } a + c = 0$$

a solution is $a = 1, b = -1, c = -1$ *MI*
AI

THEN

π_3 has equation $x - y - z = d$ *(MI)*
as it goes through the origin, $d = 0$ so π_3 has equation $x - y - z = 0$ *AI*

Note: The final *(MI)AI* are independent of previous working.

Question 3 continued

METHOD 2

$$r = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + s \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix} + t \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$$

AI(AI)AIAI

[7 marks]

4. $2^{2x-2} = 2^x + 8$

(M1)

$$\frac{1}{4} 2^{2x} = 2^x + 8$$

(A1)

$$2^{2x} - 4 \times 2^x - 32 = 0$$

A1

$$(2^x - 8)(2^x + 4) = 0$$

(M1)

$$2^x = 8 \Rightarrow x = 3$$

A1

Notes: Do not award final *A1* if more than 1 solution is given.

[5 marks]


5. (a) an attempt to use either asymptotes or intercepts

(M1)

$$a = -2, b = 1, c = \frac{1}{2}$$

AIAIAI

(b)



A4

Note: Award *A1* for both asymptotes,
A1 for both intercepts,
A1, A1 for the shape of each branch, ignoring shape at $(x = -2)$.

[8 marks]

6. $(a + b) \cdot (a - b) = a \cdot a + b \cdot a - a \cdot b - b \cdot b$ *MI*
 $= a \cdot a - b \cdot b$ *AI*
 $= |a|^2 - |b|^2 = 0$ since $|a| = |b|$ *AI*
 the **diagonals** are perpendicular *RI*

Note: Accept geometric proof, awarding *MI* for recognizing OACB is a rhombus, *RI* for a clear indication that $(a + b)$ and $(a - b)$ are the diagonals, *AI* for stating that diagonals cross at right angles and *AI* for “hence dot product is zero”.

Accept solutions using components in 2 or 3 dimensions.

[4 marks]

7. $P(\text{six in first throw}) = \frac{1}{6}$ *(AI)*
 $P(\text{six in third throw}) = \frac{25}{36} \times \frac{1}{6}$ *(MI)(AI)*
 $P(\text{six in fifth throw}) = \left(\frac{25}{36}\right)^2 \times \frac{1}{6}$
 $P(\text{A obtains first six}) = \frac{1}{6} + \frac{25}{36} \times \frac{1}{6} + \left(\frac{25}{36}\right)^2 \times \frac{1}{6} + \dots$ *(MI)*
 recognizing that the common ratio is $\frac{25}{36}$ *(AI)*
 $P(\text{A obtains first six}) = \frac{\frac{1}{6}}{1 - \frac{25}{36}}$ (by summing the infinite GP) *MI*
 $= \frac{6}{11}$ *AI*

[7 marks]

8. $\sqrt{x} e^x = e\sqrt{x} \Rightarrow x = 0$ or 1 *(AI)*

attempt to find $\int y^2 dx$ *MI*

$$V_1 = \pi \int_0^1 e^2 x dx$$

$$= \pi \left[\frac{1}{2} e^2 x^2 \right]_0^1$$

$$= \frac{\pi e^2}{2} \quad \text{AI}$$

$$V_2 = \pi \int_0^1 x e^{2x} dx$$

$$= \pi \left(\left[\frac{1}{2} x e^{2x} \right]_0^1 - \int_0^1 \frac{1}{2} e^{2x} dx \right) \quad \text{MIAI}$$

Note: Award *MI* for attempt to integrate by parts.

$$= \frac{\pi e^2}{2} - \pi \left[\frac{1}{4} e^{2x} \right]_0^1$$

finding difference of volumes *MI*

volume = $V_1 - V_2$

$$= \pi \left[\frac{1}{4} e^{2x} \right]_0^1$$

$$= \frac{1}{4} \pi (e^2 - 1) \quad \text{AI}$$

[7 marks]

9. (a) $u = \frac{1}{x} \Rightarrow du = -\frac{1}{x^2} dx$ *MI*

$\Rightarrow dx = -\frac{du}{u^2}$ *AI*

$$\int_1^\alpha \frac{1}{1+x^2} dx = - \int_1^\alpha \frac{1}{1 + \left(\frac{1}{u}\right)^2} \frac{du}{u^2} \quad \text{AIMIAI}$$

Note: Award *AI* for correct integrand and *MIAI* for correct limits.

$$= \int_\alpha^1 \frac{1}{1+u^2} du \quad (\text{upon interchanging the two limits}) \quad \text{AG}$$

(b) $\arctan x \Big|_1^\alpha = \arctan u \Big|_{\frac{1}{\alpha}}^1$ *AI*

$$\arctan \alpha - \frac{\pi}{4} = \frac{\pi}{4} - \arctan \frac{1}{\alpha} \quad \text{AI}$$

$$\arctan \alpha + \arctan \frac{1}{\alpha} = \frac{\pi}{2} \quad \text{AG}$$

[7 marks]

10. EITHER

$$\text{let } y_i = x_i - 12$$

$$\bar{x} = 10 \Rightarrow \bar{y} = -2$$

MIAI

$$\sigma_x = \sigma_y = 3$$

AI

$$\frac{\sum_{i=1}^{10} y_i^2}{10} - \bar{y}^2 = 9$$

MIAI

$$\sum_{i=1}^{10} y_i^2 = 10(9 + 4) = 130$$

*AI***OR**

$$\sum_{i=1}^{10} (x_i - 12)^2 = \sum_{i=1}^{10} x_i^2 - 24 \sum_{i=1}^{10} x_i + 144 \sum_{i=1}^{10} 1$$

MIAI

$$\bar{x} = 10 \Rightarrow \sum_{i=1}^{10} x_i = 100$$

AI

$$\sigma_x = 3, \frac{\sum_{i=1}^{10} x_i^2}{10} - \bar{x}^2 = 9$$

(MI)

$$\Rightarrow \sum_{i=1}^{10} x_i^2 = 10(9 + 100)$$

AI

$$\sum_{i=1}^{10} (x_i - 12)^2 = 1090 - 2400 + 1440 = 130$$

*AI**[6 marks]*

SECTION B

11. (a) $x^2 + 5x + 4 = 0 \Rightarrow x = -1$ or $x = -4$ (MI)
 so vertical asymptotes are $x = -1$ and $x = -4$ AI
 as $x \rightarrow \infty$ then $y \rightarrow 1$ so horizontal asymptote is $y = 1$ (MI)AI

[4 marks]

- (b) $x^2 - 5x + 4 = 0 \Rightarrow x = 1$ or $x = 4$ AI
 $x = 0 \Rightarrow y = 1$ AI
 so intercepts are (1, 0), (4, 0) and (0, 1)

[2 marks]

- (c) (i) $f'(x) = \frac{(x^2 + 5x + 4)(2x - 5) - (x^2 - 5x + 4)(2x + 5)}{(x^2 + 5x + 4)^2}$ MIAIAI

$$= \frac{10x^2 - 40}{(x^2 + 5x + 4)^2} \left(= \frac{10(x-2)(x+2)}{(x^2 + 5x + 4)^2} \right) \quad \text{AI}$$

$$f'(x) = 0 \Rightarrow x = \pm 2 \quad \text{MI}$$

so the points under consideration are $(-2, -9)$ and $\left(2, -\frac{1}{9}\right)$ AIAI

looking at the sign either side of the points (or attempt to find $f''(x)$) MI

e.g. if $x = -2^-$ then $(x-2)(x+2) > 0$ and if $x = -2^+$ then $(x-2)(x+2) < 0$,
 therefore $(-2, -9)$ is a maximum AI

- (ii) e.g. if $x = 2^-$ then $(x-2)(x+2) < 0$ and if $x = 2^+$ then $(x-2)(x+2) > 0$,
 therefore $\left(2, -\frac{1}{9}\right)$ is a minimum AI


Note: Candidates may find the minimum first.

[10 marks]

continued ...

Question 11 continued

(d)



A3

Note: Award **A1** for each branch consistent with and including the features found in previous parts.

[3 marks]

(e) one

A1

[1 mark]

Total [20 marks]

12. (a) $\int_0^1 ae^{-ax} dx = 1 - \frac{1}{\sqrt{2}}$ *MIAI*
 $[-e^{-ax}]_0^1 = 1 - \frac{1}{\sqrt{2}}$ *MIAI*
 $-e^{-a} + 1 = 1 - \frac{1}{\sqrt{2}}$ *AI*

Note: Accept e^0 instead of 1.

$e^{-a} = \frac{1}{\sqrt{2}}$
 $e^a = \sqrt{2}$
 $a = \ln 2^{\frac{1}{2}}$ (accept $-a = \ln 2^{\frac{1}{2}}$) *AI*
 $a = \frac{1}{2} \ln 2$ *AG*

[6 marks]

(b) $\int_0^M ae^{-ax} dx = \frac{1}{2}$ *MIAI*
 $[-e^{-ax}]_0^M = \frac{1}{2}$ *AI*
 $-e^{-Ma} + 1 = \frac{1}{2}$
 $e^{-Ma} = \frac{1}{2}$ *AI*
 $Ma = \ln 2$
 $M = \frac{\ln 2}{a} = 2$ *AI*

[5 marks]

continued ...

Question 12 continued

(c) $P(1 < X < 3) = \int_1^3 ae^{-ax} dx$	<i>MIAI</i>
$= -e^{-3a} + e^{-a}$	<i>AI</i>

$P(X < 3 X > 1) = \frac{P(1 < X < 3)}{P(X > 1)}$	<i>MIAI</i>
$= \frac{-e^{-3a} + e^{-a}}{1 - P(X < 1)}$	<i>AI</i>
$= \frac{-e^{-3a} + e^{-a}}{\frac{1}{\sqrt{2}}}$	<i>AI</i>
$= \sqrt{2}(-e^{-3a} + e^{-a})$	
$= \sqrt{2}\left(-2^{-\frac{3}{2}} + 2^{-\frac{1}{2}}\right)$	<i>AI</i>
$= \frac{1}{2}$	<i>AI</i>

Note: Award full marks for $P(X < 3 / X > 1) = P(X < 2) = \frac{1}{2}$ or quoting properties of exponential distribution.

[9 marks]

Total [20 marks]

13. (a) $\sin (2n+1)x \cos x - \cos (2n+1)x \sin x = \sin (2n+1)x - x$
 $= \sin 2nx$

MIAI
AG

[2 marks]

(b) if $n = 1$

MI

LHS = $\cos x$

RHS = $\frac{\sin 2x}{2 \sin x} = \frac{2 \sin x \cos x}{2 \sin x} = \cos x$

MI

so LHS = RHS and the statement is true for $n = 1$

RI

assume true for $n = k$

MI

Note: Only award **MI** if the word **true** appears.
 Do **not** award **MI** for 'let $n = k$ ' only.
 Subsequent marks are independent of this **MI**.

so $\cos x + \cos 3x + \cos 5x + \dots + \cos (2k - 1)x = \frac{\sin 2kx}{2 \sin x}$

if $n = k + 1$ then

$\cos x + \cos 3x + \cos 5x + \dots + \cos (2k - 1)x + \cos (2k + 1)x$

MI

$= \frac{\sin 2kx}{2 \sin x} + \cos (2k + 1)x$

AI

$= \frac{\sin 2kx + 2 \cos (2k + 1)x \sin x}{2 \sin x}$

MI

$= \frac{\sin (2k + 1)x \cos x - \cos (2k + 1)x \sin x + 2 \cos (2k + 1)x \sin x}{2 \sin x}$

MI

$= \frac{\sin (2k + 1)x \cos x + \cos (2k + 1)x \sin x}{2 \sin x}$

AI

$= \frac{\sin (2k + 2)x}{2 \sin x}$

MI

$= \frac{\sin 2(k + 1)x}{2 \sin x}$

AI

so if true for $n = k$, then also true for $n = k + 1$

as true for $n = 1$ then true for all $n \in \mathbb{Z}^+$

RI

Note: Final **RI** is independent of previous work.

[12 marks]

continued ...

Question 13 continued

- (c) $\frac{\sin 4x}{2 \sin x} = \frac{1}{2}$ *MIAI*
 $\sin 4x = \sin x$
 $4x = x \Rightarrow x = 0$ but this is impossible
 $4x = \pi - x \Rightarrow x = \frac{\pi}{5}$ *AI*
 $4x = 2\pi + x \Rightarrow x = \frac{2\pi}{3}$ *AI*
 $4x = 3\pi - x \Rightarrow x = \frac{3\pi}{5}$ *AI*
for not including any answers outside the domain *RI*

Note: Award the first *MIAI* for correctly obtaining $8\cos^3 x - 4\cos x - 1 = 0$ or equivalent and subsequent marks as appropriate including the answers $\arccos\left(-\frac{1}{2}, \frac{1 \pm \sqrt{5}}{4}\right)$.

[6 marks]

Total [20 marks]
