



# CHEMISTRY STANDARD LEVEL PAPER 2

Friday 9 November 2012 (afternoon)

1 hour 15 minutes

Candidate session number								
0	0							

#### Examination code

8	8	1	2	_	6	1	0	5
	l		l	1		l		

### **INSTRUCTIONS TO CANDIDATES**

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- Section A: answer all questions.
- Section B: answer one question.
- Write your answers in the boxes provided.
- A calculator is required for this paper.
- A clean copy of the *Chemistry Data Booklet* is required for this paper.
- The maximum mark for this examination paper is [50 marks].

## **SECTION A**

Answer all questions. Write your answers in the boxes provided.

- 1. Two groups of students (Group A and Group B) carried out a project\* on the chemistry of some group 7 elements (the halogens) and their compounds.
  - (a) In the first part of the project, the two groups had a sample of iodine monochloride (a corrosive brown liquid) prepared for them by their teacher using the following reaction.

$$I_2(s) + Cl_2(g) \rightarrow 2ICl(l)$$

The following data were recorded.

Mass of I <sub>2</sub> (s)	10.00 g
Mass of Cl <sub>2</sub> (g)	2.24 g
Mass of ICl(l) obtained	8.60 g

(i)	State the number of significant figures for the masses of $I_2(s)$ and $ICl(l)$ .	[1]
	I <sub>2</sub> (s):	
	ICl(l):	
(ii)	The iodine used in the reaction was in excess. Determine the theoretical yield, in $g$ , of $ICl(l)$ .	[3]

<sup>\*</sup> Adapted from J Derek Woollins, (2009), *Inorganic Experiments* and Open University, (2008), *Exploring the Molecular World*.



(Question 1 continued)

Although 97.4 °C an	ing a digital thermometer, the students discovered that the reaction was othermic. State the sign of the enthalpy change of the reaction, $\Delta H$ .
Although 97.4 °C an	
Although 97.4 °C an	
Although 97.4 °C an	
97.4 °C aı	
97.4 °C aı	
	In the molar masses of ICl and $Br_2$ are very similar, the boiling point of ICl is and that of $Br_2$ is 58.8 °C. Explain the difference in these boiling points in terms termolecular forces present in each liquid.



(Question 1 continued)

(i)

(c) The students reacted ICl(l) with CsBr(s) to form a yellow solid, CsICl<sub>2</sub>(s), as one of the products. CsICl<sub>2</sub>(s) has been found to produce very pure CsCl(s) which is used in cancer treatment.

To confirm the composition of the yellow solid, Group A determined the amount of iodine in 0.2015 g of  $CsICl_2(s)$  by titrating it with 0.0500 mol dm<sup>-3</sup>  $Na_2S_2O_3(aq)$ . The following data were recorded for the titration.

Mass of $CsICl_2(s)$ taken (in $g \pm 0.0001$ )	0.2015
Initial burette reading of $0.0500 \text{ mol dm}^{-3} \text{ Na}_2\text{S}_2\text{O}_3(\text{aq})$ (in cm <sup>3</sup> ± 0.05)	1.05
Final burette reading of $0.0500 \text{ mol dm}^{-3} \text{ Na}_2\text{S}_2\text{O}_3(\text{aq})$ (in cm <sup>3</sup> ± 0.05)	25.25

Calculate the percentage of iodine by mass in CsICl<sub>2</sub>(s), correct to three

	significant figures.	[1]
(ii)	State the volume, in cm <sup>3</sup> , of $0.0500 \text{ mol dm}^{-3} \text{ Na}_2\text{S}_2\text{O}_3(\text{aq})$ used in the titration.	[1]



(Question I	' continued,

(iii)	Determine the amount, in mol, of $0.0500~\rm moldm^{-3}~Na_2S_2O_3(aq)$ added in the titration.	[1]
(iv)	The overall reaction taking place during the titration is:	
	$CsICl_2(s) + 2Na_2S_2O_3(aq) \rightarrow NaCl(aq) + Na_2S_4O_6(aq) + CsCl(aq) + NaI(aq)$	
	Calculate the amount, in mol, of iodine atoms, I, present in the sample of $CsICl_2(s)$ .	[1]
(v)	Calculate the mass of iodine, in g, present in the sample of CsICl <sub>2</sub> (s).	[1]
(vi)	Determine the percentage by mass of iodine in the sample of $CsICl_2(s)$ , correct to <b>three</b> significant figures, using your answer from $(v)$ .	[1]



(Question 1 continued)

(d)	Group B heated the yellow solid, CsICl <sub>2</sub> (s), which turned white and released a brown gas which condensed into a brown liquid.	
	Group B identified the white solid as CsCl(s). Suggest the identity of the brown liquid. [A	1]



2.

	ne the term average bond enthalpy.	[2]
	following equation represents a combustion reaction of propane, $C_3H_8(g)$ when the gen supply is limited.	
	$C_3H_8(g) + 3\frac{1}{2}O_2(g) \rightarrow 3CO(g) + 4H_2O(g)$	
(i)	Determine $\Delta H$ , the enthalpy change of the reaction, in kJ mol <sup>-1</sup> , using average bond enthalpy data from Table 10 of the Data Booklet. The bond enthalpy for the carbon-oxygen bond in carbon monoxide, CO, is 1072 kJ mol <sup>-1</sup> .	[3]



(a)	A glass container is half-filled with liquid bromine and then sealed. The system eventually reaches a dynamic equilibrium. State <b>one</b> characteristic of a system in equilibrium.					
(b)	The oxidation of sulfur dioxide is an important reaction in the Contact process used to manufacture sulfuric acid. $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g) \qquad \Delta H = -198.2 \text{ kJ}$ (i) Deduce the equilibrium constant expression, $K_c$ .					
	(i)	-			[1]	
	(ii)	Predict how each of the foll the value of $K_c$ .	owing changes affects the	position of equilibrium and	[3]	
	(ii)		owing changes affects the property of the prop	position of equilibrium and $\mathbf{Value\ of}\ \mathbf{K}_{c}$	[3]	
	(ii)				_	

(This question continues on the following page)




Increase in pressure

Addition of a catalyst


(Question 3 continued)

(c) Vanadium(V) oxide,  $V_2O_5$ , is a catalyst that can be used in the Contact process. It provides an alternative pathway for the reaction, lowering the activation energy,  $E_a$ .

(i) Define the term  $activation\ energy,\ E_{\rm a}.$  [1]



(ii) Sketch the **two** Maxwell–Boltzmann energy distribution curves for a fixed amount of gas at two different temperatures,  $T_1$  and  $T_2$  ( $T_2 > T_1$ ). Label **both** axes. [3]



## **SECTION B**

Answer one question. Write your answers in the boxes provided.

(i)	Define the terms atomic number, mass number and isotopes of an element.	[3]
	Atomic number:	
	Mass number:	
	Isotopes of an element:	
(ii)	Distinguish between the terms <i>group</i> and <i>period</i> .	[1]
(iii)	Deduce the electron arrangements of the lithium ion, Li <sup>+</sup> , and the boron atom, B.	[2]
	Li <sup>+</sup> :	
	Li':	



(iv) Naturally occurring boron exists as two isotopes with mass numbers of 10 and 11.

# (Question 4 continued)

																												o																															
		_																								_																				_													
•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	 •	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	 •	•	•	•	•	•	•	•	•	•	•	•	•	•	•	• •	•	•
•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	 •	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	 •	•	•	•	•	•	•	•	•	•	•	•	•	•	•		•	•
•	•	•	•	•	•	•	•	•	•	•	•	•	•	•		•		•	•	•	•	•	•			 •	•	•	•	•	•	•	•	•	•		•	•	•	•			•	•	•				•	•	•	•	•	•	•				•

(v) Lithium exists as two isotopes with mass numbers of 6 and 7. Deduce the number of protons, electrons and neutrons for each isotope. [2]

Mass number (A)	Number of protons	Number of electrons	Number of neutrons
6			
7			

(This question continues on the following page)



Turn over

0	uestion	4	continued	'
~		•		,

(b)

Eve	ry element has its own unique line emission spectrum.	
(i)	Distinguish between a continuous spectrum and a line spectrum.	[2]
(ii)	Draw a diagram to show the electron transitions between energy levels in a hydrogen atom that are responsible for the two series of lines in the ultraviolet and visible regions of the spectrum. Label your diagram to show <b>three</b> transitions for each series.	[4]



(Question 4 continued)

(iii) Iron is described as a transition metal. Identify the <b>two</b> most common ions of iron  (iii) Deduce the chemical formulas of lithium oxide and iron(II) oxide.  Lithium oxide:  Iron(II) oxide:	(1)	Explain why metals are good conductors of electricity and why they are malleable.	L
(iii) Deduce the chemical formulas of lithium oxide and iron(II) oxide.  Lithium oxide:			
(iii) Deduce the chemical formulas of lithium oxide and iron(II) oxide.  Lithium oxide:			
(iii) Deduce the chemical formulas of lithium oxide and iron(II) oxide.  Lithium oxide:			
(iii) Deduce the chemical formulas of lithium oxide and iron(II) oxide.  Lithium oxide:			
(iii) Deduce the chemical formulas of lithium oxide and iron(II) oxide.  Lithium oxide:			
Lithium oxide:	(ii)	Iron is described as a transition metal. Identify the <b>two</b> most common ions of iron.	
Lithium oxide:			
Lithium oxide:			
	(iii)	Deduce the chemical formulas of lithium oxide and iron(II) oxide.	_
Iron(II) oxide:		Lithium oxide:	
Iron(II) oxide:			
		Iron(II) oxide:	



Arsenic and nitrogen play a significant role in environmental chemistry. Arsenous acid,

H<sub>3</sub>AsO<sub>3</sub>, can be found in oxygen-poor (anaerobic) water, and nitrogen-containing fertilizers

5.

conta	minate water.	
(i)	Define oxidation and reduction in terms of electron loss or gain.	[1]
	Oxidation:	
	Reduction:	
(ii)	Deduce the oxidation numbers of arsenic and nitrogen in each of the following species.	[4]
	As <sub>2</sub> O <sub>3</sub> :	
	NO <sub>3</sub> -:	
	$H_3AsO_3$ :	
	$N_2O_3$ :	
(iii)	Distinguish between the terms oxidizing agent and reducing agent.	[1]



(Question 5 continued)

(iv)	In the removal of arsenic from	contaminated	groundwater,	$H_3AsO_3$	is often	first
	oxidized to arsenic acid, H <sub>3</sub> AsO	4.				

The following unbalanced redox reaction shows another method of forming H<sub>3</sub>AsO<sub>4</sub>.

$$As_2O_3(s) + NO_3^-(aq) \rightarrow H_3AsO_4(aq) + N_2O_3(aq)$$

Deduce the balanced redox equation in **acid**, and then identify both the oxidizing and reducing agents.

[3]


- (b) Nitric acid, HNO<sub>3</sub>, is strong and nitrous acid, HNO<sub>2</sub>, is weak.
  - (i) Define an *acid* according to the Brønsted–Lowry and Lewis theories. [2]

Brønsted-	-Lowry the	ory:		
Lewis the	eory:			



(Question 5 continued)

(ii)	The Lewis	(electron dot)	structure of	nitrous	acid is	given	below.
------	-----------	----------------	--------------	---------	---------	-------	--------

	$H - \overline{Q} - \overline{N} = \emptyset$	
	Identify which nitrogen-oxygen bond is the shorter.	[1]
(iii)	Deduce the approximate value of the hydrogen-oxygen-nitrogen bond angle in nitrous acid and explain your answer.	[2]
(iv)	Distinguish between a <i>strong acid</i> and a <i>weak acid</i> in terms of their dissociation in aqueous solution.	[1]



(Question 5 continued)

(v)	Ammonia, NH <sub>3</sub> , is a weak base. Deduce the Lewis (electron dot) structure of NH <sub>3</sub> . State the name of the shape of the molecule and explain why NH <sub>3</sub> is a polar molecule.	[3]
(vi)	When lime was added to a sample of soil, the pH changed from 5 to 7. Calculate the <b>factor</b> by which the hydrogen ion concentration changes.	[1]
(vii)	One common nitrogen-containing fertilizer is ammonium sulfate. State its chemical formula.	[1]




6.

Alke	nes, a	lcohols and esters are three families of organic compounds with many commercial uses.	
ı)	(i)	State <b>two</b> industrial uses of ethene.	[
	(ii)	State the meaning of the term structural isomers.	
			—



(Question 6 continued)

(iii)  $\mathbf{X}$  is an isomer of  $C_4H_8$  and has the structural formula shown below.



Apply IUPAC rules to name this isomer. Deduce the structural formulas of two other isomers of  $C_4H_8$ .


(iv)	State the balanced chemical equation for the reaction of <b>X</b> with HBr to form <b>Y</b> .	[1]

(This question continues on the following page)



Turn over

[3]

(Question 6 continued)

(v)	Y reacts with aqueous sodium hydroxide, NaOH (aq), to form an alcohol, Z. Identify whether Z is a primary, secondary or tertiary alcohol.	[1]
(vi)	Explain <b>one</b> suitable mechanism for the reaction in (v) using curly arrows to represent the movement of electron pairs.	[4]
(vii)	Deduce the structural formula of the organic product formed when <b>Z</b> is oxidized by heating under reflux with acidified potassium dichromate(VI) <b>and</b> state the name of the functional group of this organic product.	[2]



(Question 6 continued)

(i)	Draw the ester functional group.
(ii)	Determine the empirical formula of the ester, showing your working.
(iii)	The molar mass of the ester is 116.18 g mol <sup>-1</sup> . Determine its molecular formula.
()	



Please do not write on this page.

Answers written on this page will not be marked.



Please do not write on this page.

Answers written on this page will not be marked.



Please do not write on this page.

Answers written on this page will not be marked.

