


HL Paper 1


At 700 °C, the equilibrium constant, K_c , for the reaction is 1.075×10^8 .



Which relationship is always correct for the equilibrium at this temperature?

- A. $[\text{H}_2\text{S}]^2 < [\text{H}_2]^2 [\text{S}_2]$
- B. $[\text{S}_2] = 2[\text{H}_2\text{S}]$
- C. $[\text{H}_2\text{S}] < [\text{S}_2]$
- D. $[\text{H}_2\text{S}]^2 > [\text{H}_2]^2 [\text{S}_2]$


1.0 mol of $\text{N}_2(\text{g})$, 1.0 mol of $\text{H}_2(\text{g})$ and 1.0 mol of $\text{NH}_3(\text{g})$ are placed in a 1.0 dm^3 sealed flask and left to reach equilibrium. At equilibrium the concentration of $\text{N}_2(\text{g})$ is 0.8 mol dm^{-3} .



What are the equilibrium concentration of $\text{H}_2(\text{g})$ and $\text{NH}_3(\text{g})$ in mol dm^{-3} ?

	$[\text{H}_2(\text{g})] / \text{mol dm}^{-3}$	$[\text{NH}_3(\text{g})] / \text{mol dm}^{-3}$
A.	0.2	1.2
B.	0.4	1.4
C.	0.4	0.4
D.	0.8	1.2


The equation for the reaction between two gases, A and B, is:



When the reaction is at equilibrium at 600 K the concentrations of A, B, C and D are 2, 1, 3 and 2 mol dm^{-3} respectively. What is the value of the equilibrium constant at 600 K?

- A. $\frac{1}{6}$
- B. $\frac{9}{7}$
- C. 3
- D. 6


Components X and Y are mixed together and allowed to reach equilibrium. The concentrations of X, Y, W and Z in the equilibrium mixture are 4, 1, 4 and 2 mol dm^{-3} respectively.



What is the value of the equilibrium constant, K_c ?

- A. $\frac{1}{8}$
 - B. $\frac{1}{2}$
 - C. 2
 - D. 8
-

A mixture of 2.0 mol of H_2 and 2.0 mol of I_2 is allowed to reach equilibrium in the gaseous state at a certain temperature in a 1.0 dm^3 flask. At equilibrium, 3.0 mol of HI are present. What is the value of K_c for this reaction?




- A. $K_c = \frac{(3.0)^2}{(0.5)^2}$
 - B. $K_c = \frac{3.0}{(0.5)^2}$
 - C. $K_c = \frac{(3.0)^2}{(2.0)^2}$
 - D. $K_c = \frac{(0.5)^2}{(3.0)^2}$
-

What is the relationship between $\text{p}K_a$, $\text{p}K_b$ and $\text{p}K_w$ for a conjugate acid–base pair?

- A. $\text{p}K_a = \text{p}K_w + \text{p}K_b$
 - B. $\text{p}K_a = \text{p}K_w - \text{p}K_b$
 - C. $\text{p}K_a \times \text{p}K_b = \text{p}K_w$
 - D. $\frac{\text{p}K_a}{\text{p}K_b} = \text{p}K_w$
-

The indicator, HIn is used in a titration between an acid and base. Which statement about the dissociation of the indicator, HIn is correct?




colour A

colour B

- A. In a strongly alkaline solution, colour B would be observed.
- B. In a strongly acidic solution, colour B would be observed.
- C. $[\text{In}^-]$ is greater than $[\text{HIn}]$ at the equivalence point.
- D. In a weakly acidic solution colour B would be observed.


When gaseous nitrosyl chloride, NOCl (g), decomposes, the following equilibrium is established:



2.0 mol of NOCl(g) were placed in a 1.0 dm³ container and allowed to reach equilibrium. At equilibrium 1.0 mol of NOCl(g) was present. What is the value of *K_c*?

- A. 0.50
- B. 1.0
- C. 1.5
- D. 2.0

A mixture of 0.40 mol of CO (g) and 0.40 mol of H₂ (g) was placed in a 1.00 dm³ vessel. The following equilibrium was established.




At equilibrium, the mixture contained 0.25 mol of CO (g). How many moles of H₂ (g) and CH₃OH (g) were present at equilibrium?

	Equilibrium mol of H ₂	Equilibrium mol of CH ₃ OH
A.	0.25	0.15
B.	0.50	0.25
C.	0.30	0.25
D.	0.10	0.15

Which is correct for an isolated system in equilibrium?

	Gibbs free energy	Entropy
A.	maximum	maximum
B.	maximum	minimum
C.	minimum	maximum
D.	minimum	minimum

The graph shows values of Δ*G* for a reaction at different temperatures.



Which statement is correct?

- A. The standard entropy change of the reaction is negative.
 - B. The standard enthalpy change of the reaction is positive.
 - C. At higher temperatures, the reaction becomes less spontaneous.
 - D. The standard enthalpy change of the reaction is negative.
-